

# ***Pollen of plane in the air of selected cities of Poland in 2018***

**Barbara Gałusza<sup>1</sup>, Agnieszka Lipiec<sup>2</sup>, Małgorzata Puc<sup>3, 4</sup>, Alina Stacewicz<sup>3, 4</sup>, Zenon Siergiejo<sup>5</sup>, Adam Rapiejko<sup>6, 7</sup>, Krystian Tywoniuk<sup>8</sup>, Kornel Szczygielski<sup>1</sup>, Dariusz Jurkiewicz<sup>1</sup>, Grzegorz Siergiejo<sup>9</sup>, Ewa M. Świebocka<sup>9</sup>, Piotr Rapiejko<sup>1, 5</sup>**

<sup>1</sup> Department of Otolaryngology with Division of Cranio-Maxillo-Facial Surgery in Military Institute of Medicine, Warsaw, Poland

<sup>2</sup> Department of Prevention of Environmental Hazards and Allergology, Medical University of Warsaw, Poland

<sup>3</sup> Department of Botany and Nature Conservation, Faculty of Biology, University of Szczecin, Poland

<sup>4</sup> Molecular Biology and Biotechnology Centre, Faculty of Biology, University of Szczecin, Poland

<sup>5</sup> Laboratory of Respiratory Diagnostics and Bronchoscopy, Medical University of Białystok, Poland

<sup>6</sup> Allergen Research Center, Poland

<sup>7</sup> Oxford Archaeology Ltd., Oxford, England

<sup>8</sup> Department of Otolaryngology, Audiology and Phoniatics, Children's Hospital of Bydgoszcz, Poland

<sup>9</sup> The Medical University of Białystok Children's Clinical Hospital, Poland

## **Abstract:**

The course of plane pollen season in Białystok, Bydgoszcz, Drawsko Pomorskie, Olsztyn, Opole, Szczecin, Warsaw and in Zielona Góra in the year 2018 was presented. Measurements were performed by the volumetric method (Lanzoni or Burkard pollen sampler). Pollen season was defined as the period in which 98% of the annual total catch occurred. In most of the cities, the plane pollen season in 2018 started in the second decade of April. The highest airborne concentration of 87 pollen grains/m<sup>3</sup> was noted in Zielona Góra on the April 28<sup>th</sup> and in Opole (85 pollen grains/m<sup>3</sup>) on the April 29<sup>th</sup>. The high pollen count of plane has a local character resulting from the number of trees in cities. The highest risk of pollen allergy associated with the presence of airborne plane pollen was found in Zielona Góra, Opole, Warsaw and Szczecin.

**Key words:** pollen count, plane tree, allergens, 2018

In Poland, plane tree pollen allergens are rarely the cause of allergic diseases, while in the eastern part of the Mediterranean area they show high allergenicity. In Europe, plane trees are often planted in cities because they tolerate air pollution well; many are filled in parks as well as along streets and alleys [1, 2].

## **Aim**

The aim of the study was to compare the plane tree pollen concentrations in the air of Białystok, Byd-

goszcz, Drawsko Pomorskie, Olsztyn, Opole, Szczecin, Warsaw, Zielona Góra in 2018.

## **Material and method**

The measurements of the pollen concentration in the study sites were performed with the volumetric method using Lanzoni and Burkard pollen samplers. Microscopic observations were performed on preparations obtained in a 7-day cycle with assessment of 24-hour periods [3]. The results were expressed as the

number of pollen grains in 1 m<sup>3</sup> of air per day (P/m<sup>3</sup>) [3]. The total pollen count over this period was expressed by the symbol SPI (Seasonal Pollen Index).

**Results and discussion**

The first pollen grains were observed in the air of Szczecin and Zielona Gora on April 10<sup>th</sup>, and


the last ones in Opole on May 19<sup>th</sup>. The course of the pollen seasons in each city is shown in the graphs (figs 1–4).

The highest airborne concentration of 87 pollen grains/m<sup>3</sup> was noted in Zielona Gora on April 28<sup>th</sup>, in Opole (85 pollen grains/m<sup>3</sup>) on April 29<sup>th</sup>, in Szczecin (61 grains/m<sup>3</sup>) on May 3<sup>rd</sup> and in Warsaw (43 grains/m<sup>3</sup>) on May 3<sup>rd</sup>.


**Table 1.** Characteristics of plane tree pollen season in 2018.

Site	Duration of pollen season (days)	Peak value [P/m <sup>3</sup> ] and peak date	Annual pollen sum (SPI)	Number of days concentration above threshold 15 P/m <sup>3</sup>
Białystok	16.05–15.05	4 (15.05)	27	0
Bydgoszcz	14.04–10.05	21 (4.05)	186	3
Drawsko Pomorskie	19.04–10.05	8 (7.06)	57	0
Olsztyn	19.04–12.05	12 (1.05)	46	0
Opole	17.04–12.05	85 (29.04)	693	15
Szczecin	10.04–12.05	61 (3.05)	263	6
Warsaw	23.04–13.05	43 (3.05)	285	7
Zielona Gora	13.04–8.05	87 (28.04)	753	16


**Figure 1.** Plane pollen count in Białystok and Bydgoszcz.




**Figure 2.** Plane pollen count in Drawsko Pomorskie and Olsztyn.



**Figure 3.** Plane pollen count in Opole and Szczecin.



**Figure 4.** Plane pollen count in Warsaw and Zielona Gora.



Norris-Hill and Emberlin publish *Platanus* in London as a highly sensitizing taxon of local importance [4]. Research in Poland confirms that plane pollen can reach high concentrations [5, 6].

### Conclusions

In most of the cities, the plane pollen season in 2018 started in the second decade of April.

The highest daily plane pollen concentrations were recorded in Opole and Zielona Gora. The highest risk of pollen allergy associated with the presence of airborne plane pollen was found in Zielona Gora, Opole, Warsaw and Szczecin.

3. Comtois P. *Statistical analysis of aerobiological data*. In: Mandrioli P, Comtois P, Levizzani V (eds). *Methods in Aerobiology*. Pitagora Editrice Bologna, Bologna 1998, 3.
4. Norris-Hill J, Emberlin J. *Diurnal variation of pollen concentration in the air of north-central London*. *Grana* 1991, 30: 229-234.
5. Puc M. *Pylek platanu w powietrzu Szczecina (2009-2010) (in Polish)*. *Alergoprofil* 2010, 6(3): 31-32.
6. Puc M, Rapiejko P, Lipiec A et al. *Pylek platanu w powietrzu wybranych miast Polski w 2014 r. (in Polish)*. *Alergoprofil* 2014, 10(3): 54-57.

### ORCID

- B. Gałusza – ID – [orcid.org/0000-0002-9594-5108](https://orcid.org/0000-0002-9594-5108)  
 A. Lipiec – ID – [orcid.org/0000-0003-3037-2326](https://orcid.org/0000-0003-3037-2326)  
 M. Puc – ID – [orcid.org/0000-0001-6734-9352](https://orcid.org/0000-0001-6734-9352)  
 E. Kalinowska – ID – [orcid.org/0000-0003-4821-6882](https://orcid.org/0000-0003-4821-6882)  
 K. Tywoniuk – ID – [orcid.org/0000-0002-4610-0002](https://orcid.org/0000-0002-4610-0002)  
 K. Szczygielski – ID – [orcid.org/0000-0002-3717-5424](https://orcid.org/0000-0002-3717-5424)  
 P. Rapiejko – ID – [orcid.org/0000-0003-3868-0294](https://orcid.org/0000-0003-3868-0294)

### References

1. Rapiejko P. *Alergeny pyłku roślin*. *Medical Education, Warsaw* 2012 (in Polish).
2. Łukasiewicz S. *Drzewa i krzewy polecane do obsadzeń ulicznych w miastach*. In: Drozdek M (ed). *Rośliny do zadań specjalnych*. PWSZ Sulechów 2010: 328 (in Polish).

### Author's contributions:

Gałusza B: 60%; and other Authors: 3.63% each.

Conflict of interests: The authors declare that they have no competing interests.

Financial support: Does not occur.

Ethics: The contents presented in this paper are compatible with the rules the Declaration of Helsinki, EU directives and standardized requirements for medical journals. Research in Białystok, Bydgoszcz, Drawsko Pomorskie, Olsztyn, Opole, Warsaw and Zielona Gora funded by Allergen Research Center Ltd.

*Corresponding author:*

**Barbara Gałusza, MD**

Department of Otolaryngology with Division  
of Cranio-Maxillo-Facial Surgery  
in Military Institute of Medicine  
04-141 Warsaw, Szaserów 128  
e-mail: bgalusza@wim.mil.pl